

Advanced Illness and Frailty Payer Differences 2026

	BCBSM	Priority	HAP	HUMANA	MSSP	ACO REACH
Breast Cancer	66 years and older with BOTH advanced illness and frailty	66 years of age and older by the end of the measurement period, with frailty and advanced illness	66 years or older with frailty and advanced illness.	66 years of age and older who have frailty and advanced illness	66 years of age and older with at least one claim/encounter for frailty during the measurement period AND an advanced illness diagnosis during the measurement period or year prior to the measurement period.	N/A
Colonoscopy	66 years and older with BOTH advanced illness and frailty	66 years of age and older as of December 31 of the measurement year with frailty and advanced illness / dispensed dementia medication	66 years and older with frailty and advanced illness	66 years of age and older who have frailty and advanced illness	66 years of age and older with at least one claim/encounter for frailty during the measurement period AND an advanced illness diagnosis during the measurement period or year prior to the measurement period.	N/A
Controlling High Blood Pressure (CBP)	66-80 years old with BOTH advanced illness and frailty; 81 years and older with frailty alone	66-80 years of age as of December 31 of the measurement year with frailty and advanced illness / dispensed dementia medication 81 years of age and older as of December 31 of the measurement year with frailty	66-80 years with frailty and advanced illness. 81 years and older with at least two indications of frailty	66 – 80 years of age who have frailty and advanced illness 81 years of age with frailty	66-80 years of age and older with at least one claim/encounter for frailty during the measurement period AND an advanced illness diagnosis during the measurement period or year prior to the measurement period;	N/A

	BCBSM	Priority	HAP	HUMANA	MSSP	ACO REACH
					81 years of age and older with at least one claim/encounter for frailty during the measurement period.	
Glycemic Status	66 years and older with BOTH advanced illness and frailty	66 years of age and older as of December 31 of the measurement year with frailty and advanced illness / dispensed dementia medication	66 years and older with frailty and advanced illness	66 years of age and older with frailty and advanced illness	66 years of age and older with at least one claim/encounter for frailty during the measurement period AND an advanced illness diagnosis during the measurement period or year prior to the measurement period.	N/A
Kidney Health Evaluation (KED)	66-80 years old with BOTH advanced illness and frailty; 81 years and older with frailty alone	66-80 years of age as of December 31 of the measurement year with frailty and advanced illness / dispensed dementia medication 81 years of age or older as of December 31 of the measurement year with frailty	66-80 years with frailty and advanced illness. 81 years and older with at least two indications of frailty	66 -80 years who have frailty and advanced illness 81 years of age and older with frailty		N/A
Osteoporosis Management in Women that had a Fracture (OMW)	67-80 years old with BOTH advanced illness and frailty; 81 and older	67-80 years of age as of December 31 of the measurement year with frailty and advanced illness / dispensed dementia medication. 81 years of age and older as of December 31 of the	67-80 years with frailty and advanced illness. 81 years and older with at least two	Patients 67–80 years of age with both frailty (at least two indications) and advanced		N/A

	BCBSM	Priority	HAP	HUMANA	MSSP	ACO REACH
	with frailty alone	measurement year with at least two indications of frailty with different dates of service	indications of frailty	illness (at least two different dates of service or dispensed dementia medication); Patients 81 years of age with frailty		
Statin Therapy for Patients with Cardiovascular Disease (SPC)	66 years and older with BOTH advanced illness and frailty	X	66 years and older with frailty and advanced illness	X		N/A
Eye Exam For Patients with Diabetes (EED)	66 years and older with BOTH advanced illness and frailty	66 years of age and older as of December 31 of the measurement year with frailty and advanced illness / dispensed dementia medication	66 years and older with frailty and advanced illness	Patients 66 years of age and older who have frailty and advanced illness		N/A
Statin Therapy for Patients with Diabetes (SPD)	x	66 years of age and older as of December 31 of the measurement year with frailty and advanced illness / dispensed dementia medication; members must meet both frailty and advanced illness criteria to be excluded	x	x		N/A